

PRISMA Statement

To the Editor:

In their recent commentary Takkouche and Norman¹ seriously misrepresent the PRISMA guideline for reporting systematic reviews and meta-analyses. They state that PRISMA “demand[s] the prior registration of the protocol of any systematic review and meta-analysis, requiring that this protocol should be made accessible before any hands-on work is done.” No such demands are made in PRISMA. Item 5 of the checklist asks authors to “Indicate if a review protocol exists, if and where it can be accessed (such as Web address), and, if available, provide registration information including registration number.”² The rationale for this request is: “registration may reduce unnecessary duplication of the same review question, particularly in jurisdictions with limited budgets; minimize study publication bias and selective reporting bias; and provide important information for those involved in updating systematic reviews.” That request anticipated the development of SR registration; indeed an international register is now being developed.³

Likewise, the PRISMA reporting guideline makes no statements as to the need for publication of a protocol prior to the initiation of the systematic review. It is possible that, as with the advent of clinical-trial registration, more avenues will appear whereby authors can publish protocols of such reviews. The authors also misrepresent the nature of registration, suggesting that this process would involve a judgement of originality and scientific merit of the planned research; we know of no such proposal.

Takkouche and Norman¹ also state “We also dispute the view that subgroup analyses and similar approaches should be used only if they were prespecified.” Again PRISMA makes no such demands on authors. Rather, the PRISMA explanatory paper says: “Having a protocol

can help restrict the likelihood of biased post hoc decisions in review methods, such as selective outcome reporting.”² In our explanatory article about PRISMA,² we also recognized that “Authors may modify protocols during the research, and readers should not automatically consider such modifications inappropriate.” PRISMA seeks transparency and clarity: authors need to report how the systematic review was conducted to enable all readers to decide upon its merits.

Lastly, we are not as confident as Takkouche and Norman about the effectiveness of peer review. A recent systematic review⁴ indicates that the peer review process is at best marginal and at worst ineffective. The scant research into the decision-making process of editors supports the need for clear and transparent reporting.⁵

David Moher

Ottawa Methods Centre
Ottawa Hospital Research Institute
Ottawa, Ontario, Canada
dmoher@ohri.ca

Douglas G. Altman

Centre for Statistics in Medicine
University of Oxford
Oxford, United Kingdom

Alessandro Liberati

Università di Modena e Reggio Emilia
Centro Cochrane Italiano
Milan, Italy

Jennifer Tetzlaff

Ottawa Methods Centre
Ottawa Hospital Research Institute
Ottawa, Ontario, Canada

REFERENCES

1. Takkouche B, Norman G. Meta-analysis protocol registration: *Sed quis custodiet ipsos custodes?* [But Who Will Guard the Guardians?] [commentary]. *Epidemiology*. 2010; 21:614–615.
2. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med*. 2009;6:e1000100.
3. Booth A, Clarke M, Ghersi D, Moher D, Petticrew M, Stewart L. An international registry of systematic-review protocols. *Lancet*. 2010 Jul 12. [Epub ahead of print].
4. Jefferson T, Rudin M, Brodney Folse S, Daviddoff F. Editorial peer review for improving

the quality of reports of biomedical studies. *Cochrane Database Syst Rev*. 2007; MR000016.

5. Dickersin K, Ssemanda E, Mansell C, Rennie D. What do the JAMA editors say when they discuss manuscripts that they are considering for publication? Developing a schema for classifying the content of editorial discussion. *BMC Med Res Methodol*. 2007;7:44.

The authors respond:

We thank the authors¹ for their interesting comments in defense of their proposed guidelines. However, we judge their remarks to be about semantics, not substance. The authors themselves have previously stated that PRISMA advocates protocol registration.² In our view, the main issue is not whether PRISMA “demands,” “advocates,” or “requests” registration. If the guideline was adopted as standard by journal editors, the “request” would of course become a requirement, and the real question is whether this requirement makes sense. Our view, as detailed in our original commentary,³ is that it does not: this additional layer of academic bureaucracy would merely obstruct free scientific enquiry.

Bahi Takkouche

Department of Preventive Medicine
University of Santiago de Compostela
Santiago de Compostela, Spain
CIBER en Epidemiología y Salud Pública
(CIBER-ESP)
Spain
bahi.takkouche@use.es

Guy Norman

Robens Centre for Public and
Environmental Health
Faculty of Health and Medical Sciences
University of Surrey
Surrey, United Kingdom

REFERENCES

1. Moher D, Altman DG, Liberati A, Tetzlaff J. PRISMA statement. *Epidemiology*. 2010;22:128.
2. Booth A, Clarke M, Ghersi D, Moher D, Petticrew M, Stewart L. An international registry of systematic-review protocols. *Lancet*. 2010 Jul 12. [Epub ahead of print].
3. Takkouche B, Norman G. Meta-analysis protocol registration: *sed quis custodiet ipsos custodes?* [but who will guard the guardians?] [commentary]. *Epidemiology*. 2010;21:614–615.

Nonalcoholic Fatty Liver Disease and Acute Ischemic Stroke

To the Editor:

Nonalcoholic fatty liver disease is associated with a higher risk of self-reported cardiovascular disease.¹ A recent systematic review also found persons with nonalcoholic fatty liver disease to have a 13% relative increase in carotid intima-media thickness.² The risk of acute ischemic stroke in relation to inflammatory markers of nonalcoholic fatty liver disease remains unknown.

We conducted a cross-sectional records-based study at the London Health Sciences Centre in London, Ontario. We included adults aged 20 to 75 years with suspected acute stroke between January 2005 and December 2009; had all undergone diffusion-weighted magnetic resonance imaging (MRI), which is sensitive and specific for early changes of acute ischemic stroke. Persons with an acute ischemic stroke were considered cases, and those whose scan was negative served as controls. Study entry required that a patient have had serum alanine aminotransferase and aspartate aminotransferase concentrations measured within 90 days before, or within 72 hours after, the MRI. Patients with evidence of intracranial hemorrhage or malignancy on MRI were excluded.

The primary study outcome was biochemical evidence of inflammatory nonalcoholic fatty liver disease, defined as an elevated serum alanine aminotransferase concentration ≥ 95 th percentile among the controls.

The study was done in accordance with a research protocol approved through the Research Ethics Boards of the London Health Sciences Centre.

We included 103 cases with, and 200 controls without, acute stroke, confirmed by diffusion-weighted im-

TABLE. Risk of Stroke Associated With Biochemical Markers of Inflammatory Nonalcoholic Fatty Liver Disease

Criterion for Biochemical Presence of Inflammatory Nonalcoholic Fatty Liver Disease	Prevalence of Biochemical Marker		Crude OR (95% CI)	Adjusted ^a OR (95% CI)
	Cases No. (%)	Controls No. (%)		
Elevated alanine aminotransferase ^b				
Absent ^c	84 (82)	188 (94)	1.0	1.0
Present	19 (19)	12 (6)	3.5 (1.7–7.6)	3.3 (1.3–8.4)
Elevated aspartate aminotransferase ^b				
Absent ^c	90 (87)	190 (95)	1.0	1.0
Present	13 (13)	10 (5)	2.7 (1.2–6.5)	3.6 (1.1–11.0)
Elevated aspartate aminotransferase:alanine aminotransferase ratio <2.0				
Absent ^c	79 (77)	183 (92)	1.0	1.0
Present	24 (23)	17 (9)	3.3 (1.7–6.4)	3.1 (1.4–7.0)

^aAdjusted for age (continuous in years), sex, current smoking, current heavy ethanol intake, history of hypertension, or current use of an antihypertensive agent, atrial fibrillation, LDL-cholesterol concentration (continuous in mmol/L), serum glucose concentration (continuous in mmol/L), and serum creatinine concentration (continuous in μ mol/L).

^bAn alanine aminotransferase (40.0 U/L) or aspartate aminotransferase (37.5 U/L) concentration ≥ 95 th percentile among controls.

^cReference category.

aging MRI. Details of participant characteristics are listed in the eAppendix (<http://links.lww.com/EDE/A443>).

Transaminases were measured within a median of 2 days (cases) and 17 days (controls) of the diffusion-weighted imaging MRI. The adjusted odds ratio (OR) for acute stroke in the presence of an elevated alanine aminotransferase was 3.3 (95% confidence interval [CI] = 1.3–8.4) (Table). Similar elevations were observed for aspartate aminotransferase concentration, as well as for alanine aminotransferase or aspartate aminotransferase in conjunction with an aspartate aminotransferase:alanine aminotransferase ratio under 2.0 (Table).

A study strength was the inclusion of both cases and controls who underwent sensitive diffusion-weighted MRI imaging for the assessment of acute stroke. This likely reduced the presence of diagnostic suspicion or referral bias,³ and correctly assigned persons with acute stroke as cases and those without acute stroke as controls. Second, we adjusted for a number of stroke risk factors, but not serum triglycerides or hepatic fat on imaging studies.^{1,4}

Others have observed nonalcoholic fatty liver disease to predict the future risk

of cardiovascular disease independent of metabolic syndrome.^{1,4} Among 1221 healthy Japanese men and women, the adjusted OR was 4.1 (95% CI = 1.6–11), but there were only 22 cardiovascular events, of which 12 were self-reported ischemic strokes.⁴ In a second study of 248 diabetic cases with cardiovascular disease, including just 29 nonfatal ischemic strokes, the associated OR between nonalcoholic fatty liver disease and cardiovascular disease was 1.5 (95% CI = 1.1–1.7).¹ Our study, which focused exclusively on acute ischemic stroke, complements these findings.

Confirmatory data are needed regarding whether nonalcoholic fatty liver disease is an independent risk factor for ischemic stroke. If so, then measurement of alanine aminotransferase or aspartate aminotransferase—both of which are readily available and inexpensive—could be considered along with traditional stroke risk factors such as serum glucose, lipids, and blood pressure.

Ivan Ying

Department of Medicine
Schulich School of Medicine and Dentistry
London Health Sciences Center
University of Western Ontario
London, Ontario

SDC Supplemental digital content is available through direct URL citations in the HTML and PDF versions of this article (www.epidem.com).

Gustavo Saposnik

Division of Neurology
Department of Medicine
St. Michael's Hospital
University of Toronto
Toronto, Ontario

Marian J. Vermeulen

Institute for Clinical Evaluative Sciences
University of Toronto
Toronto, Ontario

Andrew Leung

Department of Medical Imaging
Schulich School of Medicine and Dentistry
London Health Sciences Centre
University of Western Ontario
London, Ontario

Joel G. Ray

Departments of Medicine, Obstetrics and
Gynecology
Health Policy Management and Evaluation
St. Michael's Hospital
University of Toronto
Toronto, Ontario
rayj@smh.toronto.on.ca

REFERENCES

1. Targher G, Bertolini L, Poli F, et al. Nonalcoholic fatty liver disease and risk of future cardiovascular events among type 2 diabetic patients. *Eur Rev Med Pharmacol Sci.* 2005;9:269–271.
2. Sookian S, Pirola CJ. Non-alcoholic fatty liver disease is strongly associated with carotid atherosclerosis: a systematic review. *J Hepatol.* 2008;49:600–607.
3. Vandenbroucke JP, Cannegieter SC, Rosendaal FR. Travel and venous thrombosis: an exercise in thinking about bias. *Ann Intern Med.* 2009; 151:212–213.
4. Hamaguchi M, Kojima T, Takeda N, et al. Nonalcoholic fatty liver disease is a novel predictor of cardiovascular disease. *World J Gastroenterol.* 2007;13:1579–1584.

Risk of Parotid Malignant Tumors in Israel (1970–2006)

To the Editors:

Major salivary gland cancers are rare and with few known risk factors. We have analyzed the incidence and epidemiologic features of major salivary gland cancer (ICD.C07–C09¹) using data from the Israel National Cancer Registry, 1970–2006.

The annual incidence of major salivary gland cancer was 0.8 cases per

100,000. The most common subtype of salivary gland tumor was salivary gland carcinoma (60% of all cases) followed by squamous cell carcinoma (18%) and lymphoma (10%). Of 1924 cases (1039 men, 885 women), 60% were in the parotid gland, 7% in the submandibular gland, and fewer than 1% in the sublingual gland; the remaining 33% were recorded as overlapping or not otherwise specified (ICD.C 08.8, 08.9).¹ Our results for sex, age, site, tumor type, and rate per year are similar to those reported in the literature. Comparing the parotid gland tumors with all other salivary gland locations, there were no important differences in risk by sex, age, or ethnicity.

Trends for parotid gland cancers have seldom been reported, and have not been compared with other types of salivary gland tumors. The total number of parotid gland cancers in Israel increased 4-fold from 1970 to 2006 (from 16 to 64 cases per year), whereas other major salivary gland cancers remained stable (Fig. 1). The steepest increase occurred after 2001, with an average of 37 cases of parotid gland cancer annually before that date and 61 cases per year subsequently. The distribution of cases by age, sex, or tumor type did not change substantially during this time. The pop-

ulation of Israel increased 2.1-fold from 1970 to 2001, but only 1.1-fold from 2001 to 2006.² Population growth could not, therefore, explain the increased number of parotid gland cancers.

The proportion of major salivary gland cancers “not otherwise specified” declined from 36% before 2001 to 13% during 2001–2006. This shift may have contributed to changes over time in specific diagnoses. Nevertheless, even if we assume that all cases not-specified were parotid gland cancers, there remains a substantial increase of parotid gland cancers with time (72 cases per year after 2001 compared with 44 cases per year before).

Published data have suggested possible etiologies for major salivary gland cancers, including exposure to radiation (therapeutic and diagnostic), aging, and other factors.^{3–5} The marked increase of parotid gland cancers in the last decade may reflect differences in an external risk factor specific to the parotid region. The INTERPHONE study⁶ has investigated parotid gland tumors and other tumors that originate in tissues most exposed and proximal to cellular phones. Israelis are heavy users of cellular phones, with a 6-fold increase in usage (by minutes) from 1997 to 2006.^{7,8} Investigators have previously suggested a potential association of




FIGURE. For trend analyses, we added regression lines and calculated R^2 values. Parotid gland cancer: $R^2 = 0.83$; Submandibular gland cancer: $R^2 = 0.36$; Sublingual gland cancer: $R^2 = 0.02$.

parotid gland tumors with cell phones in an Israeli case-control study.⁷

Data on individual exposures to cell phones are not available in the registry data, and no causal association with parotid malignant tumors can be ascertained from these ecologic data. Further research is required to investigate the spectrum of possible etiological factors.

Rakefet Czerninski

Department of Oral Medicine
Hebrew University-Hadassah School of
Dental Medicine
Jerusalem, Israel
rakefetc@hadassah.org.il

Avi Zini

Harold D. Sgan-Cohen

Department of Community Dentistry
Hadassah School of Dental Medicine
Hebrew University
Jerusalem, Israel

REFERENCES

- World Health Organization. *International Classification of Diseases for Oncology, 3rd ed (ICD-0-3)*. Available at: <http://www.who.int/classifications/icd/adaptations/oncology/en/index.html>. Accessed July 27, 2010.
- Israel Central Bureau of Statistics. Available at: http://www.cbs.gov.il/reader/?Mival=cw_usr_view_Folder&ID=141. Accessed July 27, 2010.
- Zheng T, Holford TR, Chen Y, et al. Are cancers of the salivary gland increasing? Experience from Connecticut, USA. *Int J Epidemiol*. 1997;26:264–271.
- Horn-Ross PL, Ljung BM, Morrow M. Environmental factors and the risk of salivary gland cancer. *Epidemiology*. 1997;8:414–419.
- Eveson JW, Auclair P, Gnepp DR, El-Naggari AK. Tumors of the salivary glands. In: Barnes L, Eveson J, Reichart P, Sidransky D, eds. *Pathology and Genetics of Tumours of the Head and Neck*. Lyon: IARC press; 2005:209–222.
- Cardis E, Richardson L, Deltour I, et al. The INTERPHONE study: design, epidemiological methods, and description of the study population. *Eur J Epidemiol*. 2007;22:647–664.
- Sadetzki S, Chetrit A, Jarus-Hakak A, et al. Cellular phone use and risk of benign and malignant parotid gland tumors—a nationwide case-control study. *Am J Epidemiol*. 2008;167:457–467.
- Israel Government Commission Report for Communication Policy Recommendations, Ministry of Communication, 2008 [in Hebrew]. Available at: <http://www.moc.gov.il/new/documents/gronau2008/gronau1.pdf>. Accessed July 27, 2010.

On the Link Between Sufficient-cause Model and Potential-outcome Model

To the Editors:

The sufficient-cause model and the potential-outcome model have now become cornerstones for causal thinking in epidemiology. The link between these 2 models has been discussed.^{1–3} We present details of one specific situation to elucidate these 2 different causal approaches. We demonstrate a link between these 2 models under 2 binary causes X_1 and X_2 and a binary outcome D .

We let $D_{x_1x_2}(\omega)$ denote the potential outcomes for individual ω if, possibly contrary to fact, there had been interventions to set $X_1 = x_1$ and $X_2 = x_2$. For each individual ω , there would thus be 4 possible potential outcomes $D_{11}(\omega)$, $D_{10}(\omega)$, $D_{01}(\omega)$, and $D_{00}(\omega)$, which results in 16 ($= 2^4$) possible response types.^{1,4}

We could enumerate 9 different types of sufficient causes for D along with certain background causes A_k : A_1 , A_2X_1 , A_3X_2 , $A_4\bar{X}_1$, $A_5\bar{X}_2$, $A_6X_1X_2$, $A_7\bar{X}_1\bar{X}_2$, $A_8\bar{X}_1X_2$, and $A_9\bar{X}_1\bar{X}_2$, where we let \bar{X}_i denote the complement of X_i .¹ Here, the symbol A_k denotes a set of all components or factors other than the presence of X_1 , \bar{X}_1 , X_2 , and \bar{X}_2 , that may be required for a particular mechanism to operate. An individual is at risk for sufficient cause k if A_k is present. We can thus enumerate 512 ($= 2^9$) possible risk-status patterns for the sufficient causes.²

We show in the eTable (<http://links.lww.com/EDE/A440>) a complete enumeration of the 16 response types and the 512 risk-status patterns for the sufficient causes. Although a particular risk-status pattern in individual ω suffices to fix a response type, the converse is not true.^{2,4,5} Indeed, potential outcomes are

quantities specific to individuals, whereas the sufficient-cause model refers to mechanisms. Nonetheless, response types 7, 8, 10, 12, 14, 15, and 16 correspond to a unique risk-status pattern.⁵

In some cases, the effects of X_i may be in the same direction for all individuals. We say that X_1 and X_2 have positive monotonic effects on D if $D_{x_1x_2}(\omega)$ is non-decreasing in x_1 and x_2 for all individuals ω , ie, $D_{x_1x_2}(\omega) \geq D_{x'_1x'_2}(\omega)$ for $\forall \omega$ whenever $x_1 \geq x'_1$ and $x_2 \geq x'_2$.⁵ Under the assumption of positive monotonic effect of X_1 and X_2 , the individuals of response types 3, 5, 7, and 9 through 15 are excluded; and individuals of response types 1, 2, 4, 6, 8, and 16 may remain. These individuals can have, at the maximum, 446 risk-status patterns for the sufficient causes, including $A_4\bar{X}_1$, $A_5\bar{X}_2$, $A_7\bar{X}_1\bar{X}_2$, $A_8\bar{X}_1X_2$, and $A_9\bar{X}_1\bar{X}_2$. By contrast, we will say that the assumption of no preventive action of X_1 and X_2 holds if sufficient causes $A_4\bar{X}_1$, $A_5\bar{X}_2$, $A_7\bar{X}_1\bar{X}_2$, $A_8\bar{X}_1X_2$, and $A_9\bar{X}_1\bar{X}_2$ are all absent so that neither \bar{X}_1 nor \bar{X}_2 acts in a sufficient cause.⁴ These remaining individuals can have, at the maximum, 16 risk-status patterns for the sufficient causes. Therefore, the absence of \bar{X}_1 and \bar{X}_2 in any sufficient cause is stronger than the assumption of positive monotonic effects of both X_1 and X_2 . Thus, the concepts of “positive monotonic effect” and “no preventive action” should be clearly distinguished; the former is defined in the potential-outcome framework, while the latter is defined in the sufficient-component cause framework. Recently, VanderWeele⁶ distinguished these concepts by using the terms “counterfactual monotonicity” and “sufficient-cause monotonicity,” respectively.

Consideration of the correspondence between the 2 models should allow greater insight to facilitate use of each model in the appropriate contexts, clarifying the strengths of each model. Our explication would also facilitate understanding of the recent findings on the identifiability of particular sufficient causes and response types.^{5,7,8} As the

SDC Supplemental digital content is available through direct URL citations in the HTML and PDF versions of this article (www.epidem.com).

duality between the 2 models shows, the different approaches to causality provide complementary perspectives, and can be employed together to improve causal interpretations.

ACKNOWLEDGMENTS

We thank Tyler J. VanderWeele for his helpful comments on the earlier version of this letter.

Etsuji Suzuki

Department of Epidemiology
Okayama University Graduate School of
Medicine
Dentistry and Pharmaceutical Sciences
Okayama, Japan
etsuji-s@cc.okayama-u.ac.jp

Eiji Yamamoto

Department of Information Science
Faculty of Informatics
Okayama University of Science
Okayama, Japan

Toshihide Tsuda

Department of Environmental
Epidemiology
Okayama University Graduate School of
Environmental Science
Okayama, Japan

REFERENCES

- Greenland S, Poole C. Invariants and non-invariants in the concept of interdependent effects. *Scand J Work Environ Health*. 1988;14: 125–129.
- Flanders WD. On the relationship of sufficient component cause models with potential outcome (counterfactual) models. *Eur J Epidemiol*. 2006;21:847–853.
- VanderWeele TJ, Hernán MA. From counterfactuals to sufficient component causes and vice versa. *Eur J Epidemiol*. 2006;21:855–858.
- Greenland S, Lash TL, Rothman KJ. Concepts of interaction. In: Rothman KJ, Greenland S, Lash TL, eds. *Modern Epidemiology*. 3rd ed. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2008:71–83.
- VanderWeele TJ, Robins JM. The identification of synergism in the sufficient-component-cause framework. *Epidemiology*. 2007;18:329–339.
- VanderWeele TJ. Attributable fractions for sufficient cause interactions. *Int J Biostat*. 2010; 6:5. doi:10.2202/1557-4679.1202
- VanderWeele TJ, Robins JM. Empirical and counterfactual conditions for sufficient cause interactions. *Biometrika*. 2008;95:49–61.
- VanderWeele TJ. Epistatic interactions. *Stat Appl Genet Mol Biol*. 2010;9:1. doi:10.2202/1544-6115.1517

Color-coded Audio Computer-assisted Self-interview for Low-literacy Populations

To the Editor:

Self-reported measures are widely used in epidemiologic research; however, the validity of such reports may be questionable when studying sensitive topics. Self-administration of surveys removes interviewer effects and may reduce social desirability bias, but at the cost of potential subject confusion or data-entry errors. Audio computer-assisted self-interview (ACASI) systems provide for a high level of confidentiality, while removing literacy barriers.

The power structure of prisons combines with a history of abusive research involving prisoners¹ to create an environment where pressure to misreport may be especially high. Rates of illiteracy among prisoners are high,² limiting the utility of self-administered paper surveys. We examined the performance of a color-coded ACASI system for gathering sensitive data from male prisoners.

The participants completed a questionnaire regarding their adverse childhood experiences exposure as part of a survey of prisoner tobacco use.³ The Ohio State University Institutional Review Board and Ohio Department of Rehabilitation and Corrections approved all procedures, and all subjects gave informed consent to participate. The survey covered 7 categories, including physical, emotional, and sexual abuse.⁴ We collected data on a laptop computer-based, color-coded ACASI system, programmed using Visual Basic for Applications within Microsoft Access 2000 (Microsoft Corporation, Redmond, WA). The system displayed the question and responses onscreen while partici-

pants heard an audio recording of the materials over headphones. For example: “How often did a parent, step-parent, or adult living in your home swear at you, insult you, or put you down? If never, press red. If once or twice, press orange. If sometimes, press yellow. If often, press green. If very often, press blue. To skip this question, press black. To hear it again, press white.”

Participants entered their responses using a customized data entry device with 7 color-coded keys, built by modifying a number pad (Figure). The program had participants confirm each answer. For example, “You chose ‘never.’ If this is correct, press black. To change your answer, press white.”

The interviewer remained present during a sample question to adjust volume settings and answer any participant questions, but moved out of sight during survey administration. Following confirmation of the final response, the program instructed the participant to notify the interviewer.

The questionnaire was completed by 198 of 200 eligible participants (99%). Item response rates were high among those who received the questionnaire (range: 98%–100%; median = 100%). Complete responses were received from 191 men (96%); 2 men (1%) declined to answer 1 question but provided sufficient information to determine exposure to adverse childhood events for all categories, and 5 men (3%) declined to answer one or more questions in a manner that made it impossible to definitively determine their exposure.

Nearly all participants were willing and able to answer questions using the color-coded ACASI system, regardless of their prior experience with computers or comfort with reading. Intake data, collected by the Ohio Department of Rehabilitation and Correction from “self admissions, social and criminal history records,” identified physical and abuse in 8% of male inmates and sexual abuse in 5%.⁵ When using the ACASI system, participants in the current study

Supported by the Centers for Disease Control and Prevention (1R36DP001167–01) and National Cancer Institute at the National Institutes of Health (R25-CA117865–01).



FIGURE. Customized data entry keypad with color-coded keys, created by modifying a USB number pad. Top row: red, orange, yellow, green, blue. Bottom row: white, black.

reported about 3 times the prevalence of both physical (24%) and sexual (16%) abuse, possibly indicating a higher degree of comfort in sharing sensitive information when using the confidential system.

To promote dissemination of ACASI systems, the computer code used to create the system is available, for free, through the author's website for use or modification by other researchers

(<http://publichealthresearch.org/FreeCASI>), as are instructions for building a color-coded keypad. Color-coded ACASI systems, such as the one described in the present study, offer an effective means of gathering sensitive data, even when participants have difficulty reading or using computers.

ACKNOWLEDGMENTS

Development of the project was supported by the Behavioral Coopera-

tive Oncology Group of the Mary Margaret Walther Program, Walther Cancer Institute, Indianapolis, IN.

Ross M. Kauffman

College of Public Health
The Ohio State University
Columbus, OH
rmkauffm@iupui.edu

Ryan D. Kauffman

Oakhill Medical Associates
West Liberty, OH

REFERENCES

1. Kalmbach KC, Lyons PM. Ethical and legal standards for research in prisons. *Behav Sci Law.* 2003;21:671–686.
2. Linacre JM. The prison literacy problem. *Rasch Measurement Transactions.* 1996;10:473–474.
3. Kauffman RM, Ferketich AK, Murray DM, Bellair PE, Wewers ME. Measuring tobacco use in a prison population. *Nicotine Tob Res.* 2010;12:582–588.
4. Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med.* 1998; 14:245–258.
5. Bates J, Gonzalez C, Muncy V. Ohio Department of Rehabilitation and Correction: 2007 Intake Study. Available at: <http://www.drc.state.oh.us/web/Reports/intake/Intake%202007.pdf>. Accessed August 4, 2010.